

**Created:** September 2006  
**Reviewed:** August 2023  
**Revised:** August 2023

---

## DVT Prophylaxis in the Adult Trauma Patient

---

**Purpose:** To provide guidelines for DVT prophylaxis in the adult trauma patient

**Definitions:** Adult is defined as a patient greater than or equal to 16 years of age

**Guidelines:**

A. Chemical DVT Prophylaxis

- a. Should be initiated within the first 24 hours after patient arrival
- b. Contraindications may include
  - i. Patients with active bleeding, coagulopathy, or anticoagulation at time of admission and not reversed
  - ii. Patients who are ambulatory with anticipated discharge within 24 hours of arrival
  - iii. Patients who are ambulatory (BMAT 4)
  - iv. Patients with intracranial hemorrhage, refer to flowchart marked Attachment A
- c. If the patient is going to the operating room within 24 hours of arrival, may hold chemical DVT prophylaxis until after the surgery. The chemical DVT prophylaxis should be initiated within 24 hours post-op unless contraindicated.

B. Chemical DVT Prophylaxis


- a. Trauma patients with multiple injuries:
  - i. Use anticoagulant (i.e. Lovenox/Heparin) for chemical DVT prophylaxis
- b. Trauma patients with intracranial hemorrhage
  - i. See Parkland Protocol flowsheet marked Attachment A
- c. Trauma patients with Solid Organ Injuries
  - i. Consider chemical DVT prophylaxis within 24 hours of admission when Hgb stabilizes
- d. Trauma patients with isolated orthopedic injuries:
  - i. ASA 81mg bid while in the hospital and at discharge as appropriate.
  - ii. May use other medications per physician discretion (i.e. Eliquis, Lovenox, Heparin)
- e. Trauma patients with known history of VTE or clotting risk factor (Factor V Leiden, Protein C or S Deficiency) with pelvic to calcaneus fractures
  - i. Recommend an anticoagulant for 35 days post discharge

- f. Spinal Cord Injury:
  - i. Recommend chemical prophylaxis within 72 hours of injury. Consider chemical prophylaxis for up to 35 days post discharge for patients with major spine and spinal cord injury.
  - ii. To be managed by Neurosurgical specialist as appropriate
  
- C. Vena cava filters should be reserved for those patients at extremely high risk of complication from DVT chemoprophylaxis for a prolonged period and cannot be clinically anticoagulated

[Intentionally blank]

# Attachment A

## The Parkland Protocol



If Nsg signs off case, TS can initiate prophylactic enoxaparin.

If positive for DVT/PE, consider placement of IVC filter if therapeutic anticoagulation is contraindicated.

## References:

- G. H., Akl, E. A., Crowther, M., Gutterman, D. D., & Schuünemann, H. J. (2012). Executive summary: antithrombotic therapy and prevention of thrombosis: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*, 141(2), 7S-47S. BTF guidelines, 2017
- Brain Trauma Foundation guidelines, 2017
- Abdel-Aziz, H; Dunham, C. M., Malik, R. J., and Hileman, B. M. (2015). Timing for deep vein thrombosis chemoprophylaxis in traumatic brain injury: An evidenced based review. *Critical Care*, 19(96), 1-10. Doi: 10.1186/s13054-015-0814z
- Pastorek, R. A., Cripps, M. W., Bernstein, I. H., Scott, W. H., Madden, C. J., Rickert, K. L., Wolf, S. E., & Phelan, H. A. (2014). The Parkland Protocol's Modified Berne-Norwood criteria predict two tiers of risk for traumatic brain injury progression. *Journal of Neurotrauma*, 31, 1737-1743. DOI: 10.1089/neu.2014.3366
- Phelan, H. A., Eastman, A. L., Madden, C. J., Aldy, K., Berne, J. D., Norwood, S. H., Scott, W. W.,... Minei, J. P. (2012). TBI risk stratification at presentation: A prospective study of the incident and timing of radiographic worsening in the Parkland Protocol. *Journal of Acute Care Surgery*, 73, 12-127. DOI: 10.1097/TA.0b013e3182606327
- Hecht, Jason P. PharmD, BCPS, BCCCP; Han, Emily J. PharmD; Cain-Nielsen, Anne H. MS; Scott, John W. MD, MPH; Hemmila, Mark R. MD; Wahl, Wendy L. MD Association of timing of initiation of pharmacologic venous thromboembolism prophylaxis with outcomes in trauma patients, *Journal of Trauma and Acute Care Surgery*: January 2021 - Volume 90 - Issue 1 - p 54-63 doi: 10.1097/TA.0000000000002912
- Yorkgitis, Brian, PA-C, DO, Berndtson, Allison, MD, FACS, Cross, Alisa, Kennedy, Ryan, et al. (2022). American Association for the Surgery of Trauma/American College of Surgeons-Committee on Trauma Clinical Protocol for inpatient venous thromboembolism prophylaxis after trauma. *Journal of Trauma and Acute Care Surgery*, 92, 597-604. <https://doi.org/10.1097/TA.00000000000003475>
- Pai, M., Douketis, J. D., Mandel, J., & Collins, K. A. (2020). Prevention of venous thromboembolism in adult orthopedic surgical patients. *Up-to-date contents*.
- Yorkgitis BK, Berndtson AE, Cross A, Kennedy R, Kochuba MP, Tignanelli C, Tominaga GT, Jacobs DG, Marx WH, Ashley DW, Ley EJ, Napolitano L, Costantini TW. American Association for the Surgery of Trauma/American College of Surgeons-Committee on Trauma Clinical Protocol for inpatient venous thromboembolism prophylaxis after trauma. *J Trauma Acute Care Surg*. 2022 Mar 1;92(3):597-604. doi: 10.1097/TA.00000000000003475. PMID: 34797813
- 2022 News – Landmark Clinical Study Finds Aspirin as Effective as Commonly Used Blood Thinners to Prevent Life-Threatening Blood Clots and Death After Fracture Surgery. University of Maryland School of Medicine. <https://www.medschool.umaryland.edu/news/2022/Landmark-Clinical-Study-Finds-Aspirin-as-Effective-as-Commonly-Used-Blood-Thinner-to-Prevent-Life-Threatening-Blood-Clots-and-Death-After-Fracture-Surgery.html>