

PRACTICE GUIDELINE

Effective Date: 04-01-2019

Manual Reference: Deaconess Trauma Services

TITLE: NON-ACCIDENTAL TRAUMA (NAT) IN THE PEDIATRIC PATIENT

PURPOSE: To provide a guideline for evaluation and treatment of the pediatric patient with suspected non-accidental trauma (a/k/a child abuse or child maltreatment). Refer to Hospital P&P 40-03S “Abuse/Neglect/Domestic Violence” for additional information.

GUIDELINES:

1. Have a high index of suspicion for non-accidental trauma in patients who exhibit any of the following signs, symptoms or indications
 - a. History of present injury
 - i. No history or inconsistent history
 - ii. Changing history
 - iii. Unwitnessed injury
 - iv. Delay in seeking care
 - v. Mechanism of injury is implausible based on child’s developmental age
 - vi. Prior ED visit
 - vii. Domestic Violence in the home
 - viii. Premature infant (< 37 weeks)
 - ix. Low birth weight/IUGR
 - x. Chronic medical conditions
 - b. Physical exam findings
 - i. Torn frenulum
 - ii. Failure to thrive (“FTT”) (weight, length, head circumference)
 - iii. Large head in infants (consider measuring of OFC in children < 1 year)
 - iv. Any bruise in any non-ambulating child
 - v. Any bruise in a non-exploratory location (especially the TEN region – torso, ears and neck - < 4 years)
 - vi. Bruises, marks, or scars in patterns that suggest hitting with an object
 - c. Radiographic findings
 - i. Metaphyseal fractures
 - ii. Rib fractures in infants
 - iii. Any fracture in a non-ambulating infant
 - iv. An undiagnosed healing fracture
 - v. SDH and/or SAH on neuro imaging in young children, particularly in the absence of skull fracture < 1 year

2. Consult Social Work to advise of suspected non-accidental trauma. Social Work will contact Child Protective Services (CPS) and law enforcement as necessary and document accordingly in the progress note.
3. Major Areas of Evaluation
 - a. Evaluation and patient testing/imaging/procedure should be based on the mechanism of injury and/or suspicion of NAT. Avoid unnecessary invasive assessments and radiation procedures unless indicated by thorough examination.
 - b. Complete history
 - i. Document from/by whom and if it contradicts any prior story
 - ii. Include review of prior Primary Care Physician (PCP), Emergency Department (ED), and inpatient records as well as prior radiologic studies performed if available to look for sentinel injuries
4. Head to Toe Physical Exam with Particular Attention to
 - a. Growth parameters
 - b. Thorough skin exam, including scalp and hair, on undressed patient
 - c. Palpation of legs, arms, hands, feet and ribs to feel for crepitus or deformities
 - d. Complete neurological examination
 - e. Oral examination with attention to the lips, tongue, buccal mucosa, frenula, palate and teeth
 - f. Auricle exam
 - g. Genitalia examination
5. Head Imaging
 - a. Infants < 12 months of age should have a CT scan without contrast or MRI of the brain (preferred if patient has no sign of injury and normal mental status) to evaluate for intracranial injuries. This should be performed regardless of the presence or absence of neurological findings.
 - b. Children > 12 months of age should have a CT scan without contrast if there is mental status depression or any other signs of neurological injury. This may also include external signs of head injury, such as facial bruising or scalp hematoma.
 - c. If the CT scan without contrast or MRI indicates signs of trauma, MRI of the c-spine should be considered.
 - d. If there is suspicion of a skull fracture, consider ordering a CT scan with 3D reconstruction to better clarify fracture versus suture.
6. Abdominal Imaging
 - a. Any child who presents with signs/symptoms of abdominal trauma, bruising to the abdomen or torso, or an ALT/AST that is higher than twice normal should have a CT of the abdomen/pelvis with contrast.
 - b. Consider abdominal CT if urinalysis has > 10 RBCs and/or positive stool guaiac.
7. Skeletal Survey
 - a. Children < 3 years of age should have a skeletal survey to evaluate for occult fractures. When ordering a skeletal survey, be sure to include oblique x-rays of the ribs.

- b. Children > 3 years of age can have x-rays focusing on areas of concern rather than the entire skeleton.
- c. Consider getting a full skeletal survey in children > 3 years of age with developmental delays.
- d. Follow up skeletal survey should be ordered to be obtained two weeks following the suspected trauma to check for fractures that are too acute to show up on initial survey (i.e. rib fractures). This should be performed as an outpatient with follow up by child's pediatrician or family practice physician.
 - i. If patient does not have an established pediatrician or family practice provider, a referral should be made prior to patient discharge.

8. Ophthalmology Evaluation

- a. Children < 12 months of age should have an ophthalmologic evaluation to look for retinal hemorrhages. Retinal photographs should be obtained when possible.
- b. Children > 12 months of age should have an ophthalmologic evaluation when eye injuries are suspected, when head injury is suspected, and/or when there is facial bruising.
- c. Ophthalmologic examination should be obtained as soon as possible. However, the dilated eye exam should be deferred in children with head injuries pending neurosurgery clearance.

9. Lab Evaluation

- a. The following labs should be ordered routinely on all children suspected of NAT
 - i. CBC with diff and platelets
 - ii. Amylase
 - iii. Lipase
 - iv. CMP
 - v. PT/PTT/INR
 - vi. Urinalysis with microscope
 - vii. Stool for occult blood
- b. Consider a UDS/toxicology evaluation if there is clinical suspicion of exposure to substances or in children < 2 years of age with altered mental status.
- c. Consider Vitamin D 25 Hydroxy, Calcium, Phosphorus and PTH if clinically indicated.

10. Medical Photography

- a. Medical photography should be taken as soon as possible to document any skin findings at the time of presentation, since they can change rapidly.

11. Forensic Evidence

- a. Victims of severe abuse should have a toxicology evaluation.
- b. Victims of severe abuse should have a SANE evaluation if clinical concerns regarding sexual abuse or other need for forensic evidence collection.

12. Siblings

- a. All siblings or other at risk children in the home of the patients that are victims of suspected NAT should be evaluated by their PCP within 24 hours.
- b. Upon identification of other possible at risk individuals in the home of a NAT patient, the ED should consult Social Work and request Child Protective Services (CPS) be made aware of those individuals and document accordingly in the progress note.

13. Admission

- a. Admit all patients that have a clinical indication. Patients with identified traumatic injuries or who are undergoing NAT work up will be admitted to Trauma Services with consultation to the Pediatric Intensivist.
- b. Patients undergoing NAT work-up meet criteria for inpatient status.
- c. Admit patients when there is a concern about the safety of the patient, especially if there is a disagreement between the provider and CPS.

REFERENCES:

Non-Accidental Trauma (NAT) Protocol. (n.d.). Retrieved from http://www.upstate.edu/surgery/pdf/healthcare/trauma/nat_protocol_mostrecent_1.pdf
Upstate Golisano Children's Hospital, Syracuse NY. Level 1 Pediatric Trauma Center.

Boos, S. C., MD, FAAP. (n.d.). Physical child abuse: Diagnostic evaluation and management. Retrieved from https://www.uptodate.com/contents/physical-child-abuse-diagnostic-evaluation-andmanagement?search=imaginginsuspectedchildabuse&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1

Child abuse. (2018, October 05). Retrieved from <https://www.mayoclinic.org/diseases-conditions/child-abuse/symptoms-causes/syc-20370864>

Advanced Trauma Life Support (ATLS) 10th Edition

Pediatric Trauma Society; pediatrictraumasociety.org

Hospital P&P 40-03S “Abuse/Neglect/Domestic Violence”

REVIEWED DATE	REVISED DATE
APR 2019	JUL 2019
AUG 2020	
MAY 2021	