

Patient Name:

Your colonoscopy is scheduled for:

at

with Doctor:

Location: Gateway Hospital
4011 Gateway Blvd
Newburgh, IN 47630

GI Specialty Center—MOB 6
4219 Gateway Blvd
Newburgh, IN 47630

Midtown Hospital
600 Mary St.
Evansville, IN 47710

Henderson Hospital
1305 N. Elm St.
Henderson, KY 42420
Enter through emergency department

Please check in at the information desk on the first floor, **75 minutes** prior to your appointment time. **For Henderson patients, enter through the ER entrance and go to outpatient services.** If you cannot keep your appointment, please allow three-day notice for cancellation. To cancel or reschedule your appointment, please call 812-426-9545.

Arrival Time:

INSTRUCTIONS FOR PREPARING FOR YOUR PROCEDURE

If you take **blood thinning and/or weight loss medications**, please notify your GI physician as soon as possible at 812-426-9545 for instructions on when/if to stop them. See examples below:

- Aggrenox
- Eliquis
- Phentermine
- Ticlid
- Brilinta
- Invokana
- Plavix
- Wegovy
- Coumadin
- Jardiance
- Pletal
- Xarelto
- Effient
- Ozempic
- Pradaxa
- And more

7 DAYS before your procedure

Purchase the following medications seven days prior to your procedure. If you have any kidney issues or cannot afford the prep prescribed, call the office at 812-426-9545 for an alternative prep:

- ONE bottle of Miralax 238 gm (powder)
- FOUR Dulcolax 5mg tablets (Bisocodyl)
- TWO 32-ounce bottles of Gatorade or Powerade (no red, purple or orange)

5 DAYS before your procedure

1. **Stop** all NSAIDS including: Ibuprofen (Motrin, Advil), Naproxen (Aleve, Naprosyn) Mobic and similar medications. *[Acetaminophen (Tylenol) and 81mg Aspirin are ok]*
2. **Stop** Iron pills (Ferrous Sulfate, Fermalox, Niferex)
3. **Stop** all foods that contain skins, hulls, seeds and nuts (peanuts, popcorn, grapes, beans, peels of apples/potatoes, etc.)

1 DAY before your procedure

1. **Stop** all solid food
2. **Stop** all dairy products (milk, creamer, ice cream, etc.)
3. **Stop** ANY LIQUID that is **RED, ORANGE or PURPLE** (in jello, sports drinks, juice, popsicles, etc.)
4. You may continue drinking water and clear liquids including:
 - **Juices without Pulp:** apple, white grape, white cranberry, lemonade
 - **Clear Broth:** chicken, beef or vegetable
 - **Soda:** 7-up, Coke, etc. (including diet)
 - **Other:** black coffee/tea (sugar/sweetener is ok), popsicles, plain jello, sports drinks

...Continues on back page

3 PM the day before your procedure:

1. Take **TWO** Dulcolax tablets with 8 ounces of water.

6 PM the day before your procedure: Drink 1st dose of MIRALAX solution

1. Mix **ONE** 32-ounce bottle of Gatorade or Powerade with **HALF** of the Miralax powder.
2. Drink the entire mixture within two hours.

8 PM the day before your procedure:

1. Take **TWO** Dulcolax tablets with 8 ounces of water.

THE DAY OF your procedure:

7 hours before your procedure: Drink 2nd dose of MIRALAX solution

1. Mix the other 32 ounce bottle of Gatorade or Powerade with the **REST** of the Miralax.
2. Drink the entire mixture within two hours.

1. **Stop** all water/clear liquids at _____, **four hours** prior to your procedure time.
2. **Stop** all candy, mints, chewing tobacco, smoking of all kinds and chewing gum four hours prior to procedure.
3. **Stop** diabetic medications unless otherwise directed by your doctor, but bring them with you to the procedure.
4. Please take heart, lung, blood pressure and seizure medication with a small sip of water the morning of your procedure.
5. Bring your total list of medicines with you to the procedure.
6. **Bring a responsible adult to receive instructions and provide transportation who will remain at the facility throughout your stay (2-3 hrs).** You will not be permitted to drive or take public transportation alone (cab, bus, etc.) following your exam.
7. Wear something comfortable the day of the exam, and keep valuables at home.

IMPORTANT: Please follow all instructions listed on this sheet for your safety. If you do not, your procedure could be canceled and rescheduled.

1. Do not eat solid foods the day before your procedure.
2. **Do not drink clear liquids *four hours* prior to the procedure. A sip of water with important medicines is okay.**
3. Stop certain medications prior to procedure (**Phentermine for two weeks, blood thinners according to GI MD**)
4. **If you are taking Wegovy, Ozempic, Jardiance or Invokana, please notify our office for further instructions.**
5. You must have a responsible adult to receive instructions and provide your transportation the day of your procedure. Taking any public transportation alone (cab, bus, etc) without a responsible adult is not acceptable. **This adult must stay with you for the duration of the time you are at the facility (2-3 hrs).**
6. Follow instructions given by your gastroenterologist or on this instruction sheet **ONLY. DO NOT follow instructions from any other source, including the instructions on your prep bottle/box.**

Remember: It's normal to feel abdominal cramping and bloating and have multiple loose stools during the prep period. You may also experience nausea and vomiting during the prep period. We encourage you to complete ALL of the prep if possible. A successful prep is when your bowel movements are clear yellow liquid. On occasion, the colon is not completely cleaned out, which may result in re-prepping and coming back for a second procedure. If you have any questions regarding your bowel prep, please call 812-426-9545.