

GASTROENTEROLOGY EGD PREP INSTRUCTIONS

Date:	Arrive at:

YOU MUST HAVE A DRIVER WITH YOU ON THE DAY OF THE PROCEDURE, BECAUSE YOU WILL BE SEDATED.

Please inform the office if you are on any time of blood thinner so that the proper instructions can be given to you before your procedure. If you are on a blood thinner, follow these instructions:

- Hold **Aggrenox** 7 days prior to procedure once OK'd by prescribing physician.
- Hold **Plavix** at least 5 days prior to procedure once OK'd by prescribing physician.
- Hold Coumadin, Brilinta, or Aspirin 5 days prior to procedure once OK'd by prescribing physician.
- Hold **Pradaxa, Iron, Xarelto, or Eliquis** 3 days prior to procedure once OK'd by prescribing physician.

The morning of your procedure you may only have clear liquids (nothing to eat). Then nothing to drink after

<u>Clear Liquids</u>	Not Clear Liquids
Gatorade, Pedialyte, Powerade	NO RED or PURPLE items
Clear broth, bouillon	NO alcohol
Coffee or tea without milk or creamer	NO milk or non-dairy creamers
Carbonated or non-carbonated drinks	NO noodles or vegetables in soup
Kook-aid, Jell-O, Popsicles, hard candy	NO juice with pulp
Juiceno pulp	NO liquid you cannot see through